## 普诺赛<sup>®</sup> Procell system

## IGF-I/Somatamedin C/IGF-IA, Mouse, Recombinant

Cat. No.: GPCK018

产品信息

**物种** Mouse

表达宿主 E.coli

序列信息 MGPETLCGAELVDALQFVCGPRGFYFNKPTGYGSSIRRAPQTGIVDECCFRSCDLRRLEMYCA

PLKPTKAA with polyhistidine tag at the C-terminus.

cellsystem

**检索号** P05017.2

标签 His-tag at the C-terminus

**分子量** 8.61 kDa **有效期** 12 months

生物活性 Measure by its ability to induce MCF-7 cells proliferation. The ED50 for this effect is < 2

ng/mL. The specific activity of recombinant mouse IGF-I is > 5 × 10^5 IU/mg.

产品特性

内毒素 (EU/μg) < 0.1

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored

at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1 × PBS, pH 8.0.

复溶 It is recommended to reconstitute the lyophilized protein in sterile water to a

concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the

biological activity of the protein.

背景介绍

Insulin-like growth factor 1 (IGF-1), also called somatomedin C, is a protein that in humans is encoded by the IGF1 gene. IGF-1 is a hormone similar in molecular structure to insulin. It plays an important role in childhood growth and continues to have anabolic effects in adults. A synthetic analog of IGF-1, mecasermin, is used for the treatment of growth failure.

网站: <u>www.procell.com.cn</u> 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址:湖北省武汉市高新大道858号生物医药产业园三期C4栋



