Recombinant Mouse TREM2 Protein (Fc Tag)

Catalog Number: PKSM041175



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Mouse	
Mol_Mass	43.7 kDa	
Accession	Q99NH8	
Bio-activity	Not validated for activity	
Properties		

т.	
Pro	pe rtie s
110	

> 95 % as determined by reducing SDS-PAGE. **Purity**

< 1.0 EU per ug of the protein as determined by the LAL method. Endotoxin

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

This product is provided as lyophilized powder which is shipped with ice packs. Shipping

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Formulation


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Please refer to the printed manual for detailed information. Reconstitution

Data



Background

Triggering receptor expressed on myeloid cells-2 (TREM-2) is a cell surface receptor primarily expressed on macrophage s, osteoclasts, microglia and dendritic cells. TREM-2 is one member of the TREM family, inhibiting the releasing of inflammatory mediators, so it is an important in vivo anti-inflammatory receptor. TREM-2 consists of an 18 aa signal sequence, a 153 aa extracellular domain (ECD) with one V-type Ig-like domain, a 21 aa transmembrane (TM) domain, and a 35 aa cytoplasmic tail. A soluble form of TREM-2 (TREM-2b) created by alternate splicing diverges at aa 161. TREM-2 transduces intracellular signals through the adaptor DAP12. After binding of TREM-2 with ligand, the TREM-2/DAP12 (dead-cell-activated-receptor-associated protein)-mediated signal transduction pathway causes a series of intracellular protein tyrosine phosphorylation reactions and enzymatic reactions, which then activate the myeloid cells and participate T cell responses.

For Research Use Only