

Noggin/NOG (N-8His-Flag), Human, Recombinant**Cat. No. : GPCK211****产品信息**

别名	Noggin;NOG
物种	Human
表达宿主	Human Cells
序列信息	Gln28-Cys232
检索号	Q13253
分子量	23 kDa
有效期	12 months
生物活性	Measured by its ability to inhibit BMP-2-induced alkaline phosphatase production by ATDC5 mouse chondrogenic cells The ED50 for this effect is 0.2 µg/mL in the presence of 2000 ng/mL of Recombinant Human BMP 2.

产品特性

内毒素 (EU/µg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 500 mM NaCl, 2 mM EDTA, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍网站: www.procell.com.cn

电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



Noggin is a secreted homodimeric glyco Protein that is an antagonist of bone morphogenetic Proteins (BMPs). Mature Human Noggin contains an N-terminal acidic region, a central basic heparin-binding segment and a C-terminal cysteine-knot structure. Noggin is very highly conserved among vertebrates, such that mature human Noggin shares 99%, 99%, 98%, 97% and 89% aa sequence identity with mouse, rat bovine, equine and chicken Noggin, respectively. Secreted Noggin probably remains close to the cell surface due to its binding of heparin-containing proteoglycans. Noggin binds some BMPs such as BMP4 with high affinity and others such as BMP7 with lower affinity. It antagonizes BMP bioactivities by blocking epitopes on BMPs that are needed for binding to both type I and type II Receptors. Noggin is expressed in defined areas of the adult central nervous system and peripheral tissues such as lung, skeletal muscle and skin. During culture of human embryonic stem cells (hESC) or neural stem cells under certain conditions, addition of Noggin to antagonize BMP activity may allow stem cells to proliferate while maintaining their undifferentiated state, or alternatively, to differentiate into dopaminergic neurons.

