

A Reliable Research Partner in Life Science and Medicine

## Elab Fluor® 647 Anti-Human CD37 Antibody[IPO-24]

Catalog Number: E-AB-F1063M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human Host Mouse

**Isotype** Mouse IgG2b, κ

Clone No. IPO-24

Isotype Control Elab Fluor® 647 Mouse IgG2b, κ Isotype Control[MPC-11] [Product E-AB-F09812M]

Conjugation Elab Fluor® 647

**Conjugation Information** Elab Fluor<sup>®</sup> 647 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Data**



Human peripheral blood lymphocytes are stained with Elab

Fluor<sup>®</sup> 647 Anti-Human CD37 Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names CD37;Leukocyte antigen CD37;TSPAN26;Tspan-26

 Uniprot ID
 P11049

 Gene ID
 951

#### For Research Use Only

# Elabscience®

### Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

#### **Background**

CD37 is a 40-52 kD type II transmembrane protein, also known as tetraspanin-26. It is a member of the transmembrane tetraspanin family. It can interact with integrins and other transmembrane 4 superfamily members (CD53, CD81, CD82). CD37 is expressed predominantly on B cells; low expression is detected on T cells and myeloid cells. No expression is reported on NK cells and plasma cells. It is involved in regulation of T cell proliferation.