## AF/LE Purified Anti-Human CD44 Antibody[Hermes-1]

## catalog number: E-AB-F12150

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Rat
Is otype	Rat IgG2a, κ
Clone	Hermes-1
Buffer	Sterile PBS, pH 7.2. $<$ 1.0 EU per mg of the antibody as determined by the LAL method

Applications	Recommended Dilution
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Count [10 <sup>2</sup> ]	

Human peripheral blood lymphocytes were stained with 0.2

10<sup>3</sup> 10<sup>4</sup> 10<sup>5</sup> CD44 PE-H

µg AF/LE Purified Anti-Human CD44 Antibody[Hermes-1]

(Right) and 0.2 µg Rat IgG2a, κ Isotype Control (Left),

followed by PE-conjugated Goat Anti-Rat IgG Secondary

Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles. This preparation contains no preservatives, thus it should be handled
	under aseptic conditions.
Shipping	Ice bag

## Background

CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.