

Recombinant eIF4A2 Monoclonal Antibody

catalog number: **AN302068L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Rat;Mouse
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	IgG, κ
Clone	A788
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications


Applications	Recommended Dilution
WB	1:10000-1:20000
IHC	1:200-1:1000
IP	1:50-1:100

Data





Western Blot with eIF4A2 Monoclonal Antibody at dilution of 1:20000. Lane 1: 293T, Lane 2: MCF-7, Lane 3: HeLa, Lane 4: RAW264.7

Observed-MW:48 kDa
Calculated-MW:46 kDa




Immunohistochemistry of paraffin-embedded Human cerebrum using eIF4A2 Monoclonal Antibody at dilution of 1:1000.




Western Blot with eIF4A2 Monoclonal Antibody at dilution of 1:20000. Lane 1: PC-12

Observed-MW:48 kDa
Calculated-MW:46 kDa



Immunohistochemistry of paraffin-embedded Human breast cancer using eIF4A2 Monoclonal Antibody at dilution of 1:1000.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com


Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1

Immunohistochemistry of paraffin-embedded Mouse liver using eIF4A2 Monoclonal Antibody at dilution of 1:1000.

Immunohistochemistry of paraffin-embedded Rat cardiac muscle using eIF4A2 Monoclonal Antibody at dilution of 1:1000.



Immunoprecipitation analysis using anti-eIF4A2 Monoclonal Antibody. Western blot was performed from the immunoprecipitate using eIF4A2 Monoclonal Antibody at a dilution of 1:100. Lane 1: 5% Input, Lane 2: eIF4A2 Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG Isotype

Observed-MW:48 kDa

Calculated-MW:46 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Eukaryotic initiation factor 4A (eIF4A) has an essential role in the binding of mRNA to the 43S preinitiation complex when protein synthesis begins, as eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1