

FITC Anti-Human/Monkey CD14 Antibody[M5E2]

Catalog Number: E-AB-F1209C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|--------------------------------|--|
| Reactivity | Human;Rhesus |
| Host | Mouse |
| Isotype | Mouse IgG2a, κ |
| Clone No. | M5E2 |
| Isotype Control | FITC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802C] |
| Conjugation | FITC |
| Conjugation Information | FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. |

Applications

Recommended usage

| | |
|------------|---|
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |
|------------|---|

Data



Staining of normal human peripheral blood cells with FITC Anti-Human/Monkey CD14 Antibody[M5E2] (filled gray histogram) or FITC Mouse IgG2a, κ Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

| | |
|-----------------|---|
| Storage | Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |

Antigen Information

| | |
|------------------------|--|
| Alternate Names | CD14;Monocyte differentiation antigen CD14;Myeloid cell-specific leucine-rich glycoprotein |
| Uniprot ID | P08571 |

For Research Use Only

Gene ID

929

Background

CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein also known as LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens, and in the upregulation of adhesion molecules and expression of cytokines in monocytes and neutrophils.