

Recombinant Clusterin/Apolipoprotein J/Apo-J/CLU Monoclonal Antibody

catalog number: AN300046P

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human Clusterin / Apolipoprotein J / Apo-J / CLU protein
Host	Rabbit
Isotype	IgG
Clone	A1156
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications Recommended Dilution

IHC-P	1:100-1:500
--------------	-------------


Data



Immunohistochemistry of paraffin-embedded human pancreas using Clusterin / Apolipoprotein J / Apo-J / CLU Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human ovarian cancer using Clusterin / Apolipoprotein J / Apo-J / CLU Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human appendix using Clusterin / Apolipoprotein J / Apo-J / CLU Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Clusterin, also known as complement-associated protein SP-40, Complement cytolysis inhibitor, Apolipoprotein J, Testosterone-repressed prostate message 2, Aging-associated gene 4 protein, CLU and APOJ, is a secreted protein which belongs to the clusterin family. Clusterin/Apolipoprotein J/Apo-J is an enigmatic glycoprotein with a nearly ubiquitous tissue distribution and an apparent involvement in biological processes ranging from mammary gland involution to neurodegeneration in Alzheimer's disease. Its major form, a heterodimer, is secreted and present in physiological fluids, but truncated forms targeted to the nucleus have also been identified. Clusterin/Apolipoprotein J/Apo-J is a widely distributed glycoprotein with a wide range of biologic properties. A prominent and defining feature of clusterin is its marked induction in such disease states as glomerulonephritis, cystic renal disease, renal tubular injury, neurodegenerative conditions, atherosclerosis, and myocardial infarction. Upregulation of clusterin mRNA and protein levels detected in diverse disease states and in in vitro systems have led to suggestions that it functions in membrane lipid recycling, in apoptotic cell death, and as a stress-induced secreted chaperone protein, amongst others.