LRAT Polyclonal Antibody

catalog number: E-AB-19181



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human;Mouse;Rat	Human;Mouse;Rat	
Immunogen	Fusion protein of humar	Fusion protein of human LRAT	
Host	Rabbit	Rabbit	
Isotype	IgG	IgG	
Purification	Antigen affinity purifica	Antigen affinity purification	
Conjugation	Unconjugated	Unconjugated	
buffer	Phosphate buffered solu	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilu	Recommended Dilution	
WB	1:500-1:2000	1:500-1:2000	
IHC	1:50-1:200		
Data			
•	kDa 95- 72- 55- 36- 28- 17- 10- of SP20 cell lysate using LRAT	Immunohistochemistry of paraffin-embedded Human tonsil	
Polyclonal Antibody at dilution of 1:800		tissue using LRAT Polyclonal Antibody at dilution of	
Observed-MV:Refer to figures		1:55(×200)	
Calculated-MV:26 kDa			
Preparation & Storage			
Storage	Store at -20°C Valid for 1	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping	The product is shipped	The product is shipped with ice pack, upon receipt, store it immediately at the	
	temperature recommend	temperature recommended.	
Background			

The protein encoded by this gene localizes to the endoplasmic reticulum, where it catalyzes the esterification of all-transretinol into all-trans-retinyl ester. This reaction is an important step in vitamin A metabolism in the visual system. Mutations in this gene have been associated with early-onset severe retinal dystrophy and Leber congenital amaurosis 14. Alternative splicing results in multiple transcript variants.

For Research Use Only