

Recombinant Human Ezrin/EZR Protein

Catalog Number: PKSH033684

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Mol_Mass	69.4 kDa
Accession	P15311
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 10mM HEPES, pH 7.4.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ezrin is expressed in cerebral cortex, basal ganglia, hippocampus, hypophysis, and optic nerve. The N-terminus of ezrin contains a FERM domain which is further subdivided into three subdomains. The C-terminus contains a ERM domain. As a member of the ERM protein family, Ezrin serves as an intermediate between the plasma membrane and the actin cytoskeleton. It plays a key role in cell surface structure adhesion, migration, and organization. Ezrin probably involved in connections of major cytoskeletal structures to the plasma membrane. The N-terminal FERM domain strongly binds sodium-hydrogen exchanger regulatory factor (NHERF) proteins (involving long-range allostery). The C-terminal binds to actin, phosphatidylinositol bis-phosphate (PIP2) and membrane proteins like CD44 and ICAM-2. In epithelial cells, Ezrin is required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, Ezrin is required for normal macropinocytosis.

For Research Use Only