

APC Anti-Mouse CD9 Antibody[MZ3]

Catalog Number: AN01050E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a, κ
Clone No.	MZ3
Isotype Control	APC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	--

Data



Staining of C57BL/6 murine bone marrow cells with FITC Anti-Mouse Gr-1 Antibody and APC Anti-Mouse CD9 Antibody[MZ3] (left) or APC Rat IgG2a, κ (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	DRAP-27;MRP-1;p-24
Uniprot ID	P40240
Gene ID	12527

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Background

CD9 is a surface glycoprotein of the tetraspannin family. It is expressed on a variety of cells, including nerve, muscle cells and many cells of hematopoietic origin. CD9 is found to participate in forming a large molecular cell complex with other membrane proteins, such as MHC class II, CD19, CD5 and other TM4SF molecules. It is reported that CD9 is a marker of marginal zone B cells, B1 cells and plasma cells. The diverse functions of CD9 may largely depend upon its associated molecules on different cells.