

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Rat CD11b/c Antibody[OX-42]

Catalog Number: AN00651Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat Host Mouse

Isotype Mouse IgG2a, ĸ

Clone No. OX-42

Isotype Control Elab Fluor® Violet 450 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-

F09792Q1

Conjugation Elab Fluor® Violet 450

Conjugation Information Elab Fluor® Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Staining of C57BL/6 rat bone marrow cells with Elab Fluor® Violet 450 Anti-Rat CD11b/c Antibody[OX-42] (filled gray

histogram) or Elab Fluor® Violet 450 Mouse IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Rat C3bi receptor, Rat CR3 complement receptor, ITGAM, ITGAX;AN00651

P46892 **Uniprot ID**

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 25021;499271

The OX-42 antibody reacts with the CR3 complement (C3bi) receptor expressed on monocytes, granulocytes, macrophages, dendritic cells, and NK cells. This antibody appears to recognize a common epitope shared between CD11b and CD11c (integrin αM and αX chains). The OX-42 antibody precipitates three polypeptides with apparent molecular weights of 160, 103, and 95 kD, respectively. This antibody has been shown to block the formation of complement-mediated rosettes and leukocyte migration.