

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 700 Anti-Mouse CD4 Antibody[GK1.5]

Catalog Number: E-AB-F1097M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse Host Rat

**Isotype** Rat IgG2b, κ **Clone No.** GK1.5

Isotype Control Elab Fluor®700 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842M1]

Conjugation Elab Fluor® 700

**Conjugation Information** Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Staining of C57BL/6 murine splenocytes with Elab Fluor<sup>®</sup> 700 Anti-Mouse CD4 Antibody[GK1.5] (filled gray histogram)

or Elab Fluor<sup>®</sup> 700 Rat IgG2b, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4

Web: www.elabscience.cn

 Uniprot ID
 P06332

 Gene ID
 12504

## For Research Use Only



# Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

### **Background**

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, Ick.

Rev. V1.7

Web: www.elabscience.cn