Elabscience®

Rat TIMP-1 Antibody Pair Set

Catalog No.	E-KAB-0112	Applications	ELISA
Synonyms	TIMP1, CLGI, EPA, EPO, HCI		

Kit components & Storage

Title	Specifications	Storage
Rat TIMP-1 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Rat TIMP-1 Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year.
		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0112)	
		Rat TIMP-1 Capture Antibody	Rat TIMP-1 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Rat TIMP-1 protein	Recombinant Rat TIMP-1 protein
Information	Swissprot	P30120	
Product details	Reactivity	Rat	Rat
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Rat TIMP-1 in ELISAs.	

For Research Use Only

Elabscience®

Applications

Rat TIMP-1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Rat TIMP-1 Capture Antibody	
Capture			
ELISA	1:1000-1:10000	Rat TIMP-1 Detection Antibody	Optical Density
Detection		(Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction.

For Research Use Only