

AF/LE Purified Anti-Human CD58 Antibody[TS2/9.1]

catalog number: E-AB-F10680

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|---|
| Reactivity | Human |
| Immunogen | Recombinant Human CD58 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | TS2/9.1 |
| Purification | >98%, Protein A/G purified |
| Buffer | Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method. |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 µg/mL(0.5×10 ⁶ -1×10 ⁶ cells) |
|------------|---|

Data



Human peripheral blood lymphocytes were stained with 0.2 µg AF/LE Purified Anti-Human CD58 Antibody[TS2/9.1] (Right) and 0.2 µg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions. |
| Shipping | Ice bag |

Background

CD58, also known as lymphocyte function-associated antigen 3 (LFA-3) is a 45-70 kD cell surface protein that is a member of the immunoglobulin superfamily. Alternative splicing of CD58 gives rise to transmembrane and glycosylphosphatidylinositol (GPI)-anchored forms on cell surface. CD58 is expressed on both hematopoietic and non-hematopoietic cells including B cells, T cells, monocytes, erythrocytes, endothelial cells, epithelial cells, and fibroblasts. High levels are observed on memory T cells and dendritic cells. CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of its cognate ligand, CD2, on the T cell surface.

For Research Use Only