

Recombinant GAPDH Monoclonal Antibody

catalog number: AN004430L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Recombinant protein within Mouse GAPDH |
| Host | Rabbit |
| Isotype | IgG |
| Clone | 20B8 |
| Purification | Protein A/G Purification |
| Conjugation | Unconjugated |
| Buffer | PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4 |

Applications

Recommended Dilution

| | |
|------------|------------------|
| WB | 1:20000-1:200000 |
| IHC | 1:2000-1:4000 |
| IF | 1:2000-1:4000 |



Data



Western Blot with Recombinant GAPDH Monoclonal Antibody at dilution of 1:160000. Lane 1: A431 cell lysate, Lane 2: Caco-2 cell lysate, Lane 3: Jurkat cell lysate, Lane 4: K562 cell lysate, Lane 5: HeLa cell lysate, Lane 6: PC-3 cell lysate, Lane 7: Ramos cell lysate, Lane 8: TF-1 cell lysate, Lane 9: U20-S cell lysate, Lane 10: HL-60 cell lysate, Lane 11: HT-29 cell lysate, Lane 12: Raji cell lysate, Lane 13: HepG2 cell lysate

Observed-MW:36 kDa


Calculated-MW:36 kDa



Western blot with Recombinant GAPDH Monoclonal Antibody at dilution of 1:160000. lane 1: mouse heart tissue lysate, lane 2: mouse liver tissue lysate, lane 3: mouse uterus tissue lysate, lane 4: mouse lung tissue lysate, lane 5: mouse kidney tissue lysate, lane 6: mouse brain tissue lysate, lane 7: mouse stomach tissue lysate

Observed-MW:36 kDa

Calculated-MW:36 kDa




Western Blot with Recombinant GAPDH Monoclonal Antibody at dilution of 1:160000. Lane 1: C6 cell lysate, Lane 2: PC-12 cell lysate, Lane 3: Rat heart tissue lysate, Lane 4:

Rat liver tissue lysate, Lane 5: Rat spleen tissue lysate, Lane 6: Rat brain tissue lysate, Lane 7: Rat stomach tissue lysate, Lane 8: Rat testis tissue lysate, Lane 9: Rat thymus tissue lysate, Lane 10: Rat skeletal muscle tissue lysate


Observed-MW:36 kDa

Calculated-MW:36 kDa




Immunohistochemistry of paraffin-embedded rat ovary using Recombinant GAPDH Monoclonal Antibody at dilution of 1:5000.


Immunohistochemistry of paraffin-embedded human colon cancer using Recombinant GAPDH Monoclonal Antibody at dilution of 1:5000.



Immunohistochemistry of paraffin-embedded mouse ovary using Recombinant GAPDH Monoclonal Antibody at dilution of 1:5000.



Immunofluorescence analysis of U-2 OS cells using Recombinant GAPDH Monoclonal Antibody at dilution of 1:1000



Immunofluorescence analysis of NIH/3T3 cells using Recombinant GAPDH Monoclonal Antibody at dilution of 1:1000

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

GAPDH is an enzyme of 36kDa that catalyzes the sixth step of glycolysis and thus serves to break down glucose for energy and carbon molecules. Because the GAPDH gene is often stably and constitutively expressed at high levels in most tissues and cells, it is commonly used as a loading control for western blot and as a control for qPCR.