

## Recombinant Mouse NTPDase 2/ENTPD2 Protein (His Tag)

**Catalog Number: PKSM041340**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse NTPDase 2/ENTPD2 protein Cys26-Ser462, with an C-terminal His
<b>Calculated MW</b>	49.2 kDa
<b>Observed MW</b>	63-90 kDa
<b>Accession</b>	O55026
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 50mM Tris-HCl, 10mM CaCl <sub>2</sub> , 150mM NaCl, 10% Glycerol, pH 7.5.

### Data



### Background

CD39L1 protein (ENTPD2 or NTPDase2) is a member of the ecto-nucleoside triphosphate diphosphohydrolase family which the main role is termination of purinergic signaling. CD39L1 gene encodes a precursor protein with 495 amino acid residues which generates a 437 amino acid residues mature protein after processing. It is an ecto-nucleotidase that found on the surface of vascular adventitial cells and accessory vascular cells. CD39L1 is a Ca<sup>2+</sup>- and Mg<sup>2+</sup>-dependent enzyme that activates platelets by preferentially converting ATP to ADP. CD39L1 plays a role in regulating thrombosis and inflammation which is considered to be a therapeutic target for thromboregulation and the treatment of vascular inflammation. Alternative splicing of CD39L1 gene results in multiple transcript variants.

### For Research Use Only