

Recombinant Mouse Acp5 protein (His Tag)

Catalog Number: PDMM100032

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse Acp5 protein Met1-Pro327, with an C-terminal His
Calculated MW	35.9 kDa
Observed MW	38 kDa
Accession	Q05117
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse Acp5 proteins, 2µg/lane of Recombinant Mouse Acp5 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 38 KD.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Tartrate-resistant acid phosphatase (TRACP) or acid phosphatase 5, tartrate resistant (ACP5 or TRAP) is a glycosylated monomeric metalloenzyme expressed in mammals. TRACP is associated with osteoblast migration to bone resorption sites, and, once there, TRACP is believed to initiate osteoblast differentiation, activation, and proliferation. TRACP once considered to be just a histochemical marker of osteoclasts is now recognised to be a molecule of widespread occurrence with functions in both the skeleton and the immune system. Two forms of TRACP circulate in human blood, TRACP 5a derived from macrophages and dendritic cells, and TRACP-5b derived from osteoclasts. Recent data have demonstrated the utility of TRACP-5b as a marker of osteoclast number and bone resorption, and serum TRACP-5a as a marker of inflammatory conditions. TRACP is expressed by osteoclasts, macrophages, dendritic cells and a number of other cell types. It has a critical role in many biological processes including skeletal development, collagen synthesis and degradation, the mineralisation of bone, cytokine production by macrophages and dendritic cells, macrophage recruitment, dendritic cell maturation and a role in the development of Th1 responses.