Elabscience®

RBFOX3 Polyclonal Antibody

catalog number: E-AB-12595

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | | |
|---|--|--|--|
| Reactivity | Human | | |
| | ;Mouse | | |
| Immunogen | Synthetic peptide of hu | Synthetic peptide of human RBFOX3 | |
| Host | Rabbit | Rabbit | |
| Isotype | IgG | IgG | |
| Purification | Affinity purification | ••• | |
| Conjugation | Unconjugated | | |
| Buffer | Phosphate buffered sol | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. | |
| Applications | Recommended Dilutio | n | |
| WB | 1:200-1:1000 | | |
| IHC | 1:50-200 | | |
| Data | | | |
| | 100 — 70 — 55 — | | |
| | 40 35 25 | | |
| | 35 — 25 — | THC by LSBio | |
| | ^{35–} ^{25–} is of Jurkat cell using RBFOX3 | Immunohistochemistry of paraffin-embedded Human Brain | |
| Polyclonal An | 25– 25– is of Jurkat cell using RBFOX3 tibody at dilution of 1:550 | Immunohistochemistry of paraffin-embedded Human Brain using RBFOX3 Polyclonal Antibody at dilution of | |
| Polyclonal An Observed- | ^{35–} ^{25–} is of Jurkat cell using RBFOX3 tibody at dilution of 1:550 MW:Refer to figures | Immunohistochemistry of paraffin-embedded Human Brain | |
| Polyclonal An Observed- Calcul | 25– 25– is of Jurkat cell using RBFOX3 tibody at dilution of 1:550 | Immunohistochemistry of paraffin-embedded Human Brain using RBFOX3 Polyclonal Antibody at dilution of | |
| Polyclonal An Observed- Calcul Preparation & Storage | ^{35–} ^{25–} is of Jurkat cell using RBFOX3 tibody at dilution of 1:550 MW:Refer to figures ated-MW:34 kDa | Immunohistochemistry of paraffin-embedded Human Brain using RBFOX3 Polyclonal Antibody at dilution of 1:100(Elabscience Product Detected by Lifespan). | |
| Polyclonal An Observed- Calcul | 35- 25- 25- is of Jurkat cell using RBFOX3 tibody at dilution of 1:550 MW:Refer to figures ated-MW:34 kDa Store at -20°C Valid for | Immunohistochemistry of paraffin-embedded Human Brain using RBFOX3 Polyclonal Antibody at dilution of | |

Background

Immunoprecipitation and mass spectrometry of the two major NeuN species at 45–50 kDa identified both as the RNA binding protein Rbfox3 (a member of the Fox family of alternative splicing factors), confirming and extending the identification of the 45 kDa band as Rbfox3. Mapping of the anti-NeuN reactive epitopes in both R3hdm2 and Rbfox3 reveals a common proline- and glutamine-rich domain that lies at the N-terminus of the Rbfox3 protein. Nuclear Rbfox3 isoforms can also enhance the inclusion of cryptic exons in the Rbfox2 mRNA, resulting in nonsense-mediated decay of the message, thereby contributing to the negative regulation of Rbfox2 by Rbfox3 through a novel mechanism.