A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Mouse CD279/PD-1 Antibody[29F.1A12]

Catalog Number: GF11310

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse

Immunogen Recombinant Mouse CD279 protein

**Host** Rat

**Isotype** Rat IgG2a, κ **Clone** 29F.1A12

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the

LAL method.

## Applications Recommended Dilution

FCM  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### **Data**



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse CD279 gene were stained with 0.2 μg AF/LE Purified Anti-Mouse CD279 Antibody[29F.1A12] (Right) and 0.2 μg

Rat IgG2a, κ Isotype Control (Left), followed by Elab Fluor<sup>®</sup> 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

### **Preparation & Storage**

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid

freeze / thaw cycles. This preparation contains no preservatives, thus it

should be handled under as eptic conditions.

Shipping Ice bag

## **Background**

CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the CD28 family of the lg superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for prolonged survival of intratesticular islet allografts.