Recombinant Human SELENOI/EPT1 Protein (GST Tag)

Catalog Number: PKSH032407

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Description	
Species	Human
Source	E.coli-derived Human SELENOI; EPT1 protein Met 1-Pro50, with an N-terminal GST
Calculated MW	32.6 kDa
Observed MW	29 kDa
Accession	Q9C0D9
Bio-activity	Not validated for activity
Properties	
Purity	>95% as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel
	packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM EDTA,
	pH 8.0.
Data	
kDa	MK R
120	
90 60	ence
Elabs 00	Flagse
30	

> 95 % as determined by reducing SDS-PAGE.

20

14

Background

Ethanolaminephosphotransferase 1 (EPT1) is an enzyme that belongs to the CDP-Alcohol Phosphatidyltransferase Clas s-I Family. EPT1 is a Selenoprotein, which contains a Selenocysteine (Sec) residue at its active site. The Selenocysteine is encoded by the UGA codon that normally signals translation termination. The 3' UTR of Selenoprotein genes have a common stem-loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. EPT1 catalyzes Phosphatidylethanolamine biosynthesis from CDP-Ethanolamine. It plays a central role in the formation and maintenance of vesicular membranes. EPT1 is involved in the formation of Phosphatidylethanolamine via the 'Kennedy' pathway.