

Purified Anti-Human CD273 Antibody[HIQB273-2]

catalog number: AN008630P

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human
Immunogen	Recombinant Human CD273 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	HIQB273-2
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM $\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Human CD273 gene were stained with 0.2 μg Purified Anti-Human CD273 Antibody[HIQB273-2] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

T cells require a signal induced by the engagement of the T cell receptor and a "co-stimulatory" signal(s) through distinct T cell surface molecules for optimal T cell activation and tolerance. Members of the B7 superfamily of counter-receptors were identified by their ability to interact with co-stimulatory molecules found on the surface of T cells. Members of the B7 superfamily include B7-1 (CD80), B7-2 (CD86), B7-H1 (PD-L1), B7-H2 (B7RP-1), B7-H3, and PD-L2 (B7-DC).

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2