普诺赛[®] Procell system

IL-13 (C-6His), Human, Recombinant

Cat. No.: GPCK174

产品信息

物种 Human

cell system 表达宿主 Human Cells

Gly35-Asn146 序列信息 检索号 AAH96139 标签 C-6His 分子量 13.4 KDa 有效期 12 months

生物活性 Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The

ED50 for this effect is 1.5-4.5 ng/mL.

产品特性

内毒素(EU/μg) < 0.1

Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. 保存

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

复溶 Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not

> recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.

Interleukin-13 is also known as IL-13. It is a protein that in humans is encoded by the IL13 gene. Interleukin-13 is an immunoregulatory cytokine produced primarily by activated Th2 cells. It is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of proinflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergeninduced asthma but operates through mechanisms independent of IgE and eosinophils.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



