

Recombinant B7-H4 Monoclonal Antibody

catalog number: AN301876L

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse
Immunogen	Recombinant protein of human B7-H4
Host	Rabbit
Isotype	Rabbit IgG
Clone	C211498
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

WB	1:1000
IHC-P	1:100
ICC/IF	1:50
IP	1:20-1:50
FCM	1:200-1:500

Data



Western blot analysis of T-47D cell lysates using B7-H4
Antibody (AN301876L) at a dilution of 1:1000.

Observed-MW:75 kDa

Calculated-MW:31 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

B7 homolog 4 (B7-H4), also known as VTCN1, is a member of the B7 family of cell surface ligands that regulate T cell activation and immune responses. B7-H4 protein contains two extracellular Ig-like V-type domains, a transmembrane domain, and a short, two amino acid intracellular domain. The B7-H4 protein is shown to inhibit T cell activation, proliferation, and cytokine production. Although B7-H4 mRNA is widely expressed, B7-H4 protein is restricted to antigen presenting cells and B cells. The B7-H4 protein is also found in several tumor types, including ovarian cancer and breast cancer. Research studies indicate that B7-H4 protein is present on the surface of ovarian tumor cells, and that targeted inhibition of B7-H4 using recombinant antibodies restores T cell activation pathways. These studies suggest some potential therapeutic value in blocking B7-H4 function and restoring T cell function in

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.5