Recombinant Human ICAM-2/CD102 Protein (His &Fc Tag)

Catalog Number: PKSH031678

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	HEK293 Cells-derived Human ICAM-2/CD102 protein Met 1-Gln 223, with an C-terminal His & Fc
Calculated MW	50.3 kDa
Observed MW	85-95 kDa
Accession	NP_000864.2
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of PMA-stimulated HSB2 human peripheral blood acute lymphoblastic leukemia cells. When cells are added to ICAM2-coated plates (12.5 μg/mL, 100 μL/well), approximately 50 % - 60 % will adhere specifically.

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Intercellular adhesion molecule 2 (ICAM-2, CD102), belongs to the ICAM family consisting of three members identified as ligands for integrin receptors. It is a type I transmembrane glycoprotein with two Ig-like C2-type domains, and binds to the leukocyte integrins LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). As a second ligand of leukocyte function-associated antigen-1, ICAM-2 functions as a costimulatory molecule for effector cells. ICAM-2 is mainly expressed on vascular endothelial and hematopoietic cells. Interactions of ICAM-2 and the integrin receptors mediate cell adhesion in a wide range of lymphocyte, monocyte, natural killer cell, and granulocytewith other cells, and play important roles in many adhesion-dependent immune and inflammation responses, such as T cell aggregation, NK-cell cytotoxicity and migration, lymphocyte recirculation, etc. Serum levels of ICAM-2 correlated significantly with the inflammatory and course sequences of trichinosis in mice and had a similar relation with blood eosinophilia. So, estimation of ICAM-2 serum levels may prove useful in diagnosis of trichinosis recent infections, and in monitoring the prognosis and response to treatment.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com