

## Recombinant MYPT1 Monoclonal Antibody

catalog number: **AN301350L**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


|                     |                                                                 |
|---------------------|-----------------------------------------------------------------|
| <b>Reactivity</b>   | Human                                                           |
| <b>Immunogen</b>    | Recombinant Human MYPT1 protein                                 |
| <b>Host</b>         | Rabbit                                                          |
| <b>Isotype</b>      | IgG, $\kappa$                                                   |
| <b>Clone</b>        | B1117                                                           |
| <b>Purification</b> | Protein A                                                       |
| <b>Buffer</b>       | PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant. |

### Applications

### Recommended Dilution

|              |                |
|--------------|----------------|
| <b>IHC</b>   | 1:200-1:1000   |
| <b>WB</b>    | 1:2000-1:10000 |
| <b>IF</b>    | 1:200-1:1000   |
| <b>ELISA</b> | 1:5000-1:20000 |
| <b>IP</b>    | 1:50-1:200     |

### Data



Western Blot with Recombinant MYPT1 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: Daudi cell lysate.

**Observed-MW:140 kDa**

**Calculated-MW:140 kDa**

### Preparation & Storage

|                 |                                                                 |
|-----------------|-----------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag                                                         |

### Background

Myosin phosphatase target subunit 1, which is also called the myosin-binding subunit of myosin phosphatase, is one of the subunits of myosin phosphatase. Myosin phosphatase regulates the interaction of actin and myosin downstream of the guanosine triphosphatase Rho. The small guanosine triphosphatase Rho is implicated in myosin light chain (MLC) phosphorylation, which results in contraction of smooth muscle and interaction of actin and myosin in nonmuscle cells. The guanosine triphosphate (GTP)-bound, active form of RhoA (GTP.RhoA) specifically interacted with the myosin-binding subunit (MBS) of myosin phosphatase, which regulates the extent of phosphorylation of MLC. Rho-associated kinase (Rho-kinase), which is activated by GTP. RhoA, phosphorylated MBS and consequently inactivated myosin phosphatase.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.3