



A Reliable Research Partner in Life Science and Medicine

Recombinant S100A9 Monoclonal Antibody

catalog number: AN300081P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human S100A9 Protein

Host Rabbit
Isotype IgG
Clone A1191
Purification Protein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

WB 1:500-1:2000

Data



Western Blot with S100A9 Monoclonal Antibody at dilution of 1:500. Lane A: HL-60 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:15 kDa
Calculated-MW:13 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

For Research Use Only

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

S100 protein is a family of low molecular weight protein found in vertebrates characterized by two EF-hand calciumbinding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is 100% soluble in ammonium sulfate at neutral pH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from the neural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A9, also known as S100 calciumbinding protein A9, S100A9, and CAGB, is a member of the S-100 family. S100A9 is expressed by macrophages in acutely inflammed tissues and in chronic inflammation. It is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a calcium-binding protein. It has anti-microbial activity towards bacteria and fungi. The anti-microbial and proapoptotic activity of S100A9 is inhibited by zinc ions. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. S100A9 plays a role as a proinflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.

For Research Use Only