

Recombinant Human SDF2 Protein (His Tag)

Catalog Number: PKSH031339

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human SDF2 protein Met 1-Leu 211, with an C-terminal His
Calculated MW	22.7 kDa
Observed MW	22.7 kDa
Accession	NP_008854.2
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, pH 8.0, 0.5mM PMSF, 0.5mM EDTA, 0.5mM TCEP, 10% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Stromal derived factors (SDFs) are a loosely defined group of molecules that are generated by stromal cells. Two of the stromal derived factors, SDF-1 and SDF-4 belong to the chemokine family. Other SDFs, such as SDF-2 and SDF-5 are not well defined and their biological functions are less known. SDF-2 is first isolated from the mouse stromal cell line ST2 as a secretory protein. The amino acid sequence deduced from the murine clone and the human homolog are conserved more than 92 %, and the aa sequence of SDF-2 shows similarity to those of yeast dolichyl phosphate-D-mannose, protein mannosyltransferases. SDF-1 and its receptor are strongly indicated in the progression of various cancers including breast cancer. SDF-2, SDF2-L1, SDF-4, and SDF-5 are ubiquitously expressed in various cancer cell lines and SDF-2, SDF-4 and SDF-5 are expressed in mammary tissues. These SDFs have prognostic value and warrant further investigation in their biological functions and clinical value.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017