## Recombinant Rat Angiopoietin-1/ANG1 protein (His Tag)

## Catalog Number: PDER100186

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Description		
Calculated MW16.1 kDaObserved MW18 kDaAccession035460Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Species	Rat	
Observed MW18 kDaAccessionO35460Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Findotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Rat Angiopoietin-1 protein Asn304-Ala450, with an N-terminal His	
AccessionO35460Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Findotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	16.1 kDa	
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	18 kDa	
Properties   Purity > 95% as determined by reducing SDS-PAGE.   Endotoxin < 10 EU/mg of the protein as determined by the LAL method   Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.   Shipping This product is provided as lyophilized powder which is shipped with ice packs.	Accession	O35460	
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.	Bio-activity	Not validated for activity	
Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.	Properties		
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	>95% as determined by reducing SDS-PAGE.	
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method	
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
<b>Shipping</b> This product is provided as lyophilized powder which is shipped with ice packs.		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Example in the second sec	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
<b>Formulation</b> Lyophilized from a 0.2 µm intered solution in PBS with 5% Trenatose and 5%	Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%	
Mannitol.		Mannitol.	
<b>Reconstitution</b> It is recommended that sterile water be added to the vial to prepare a stock solution	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of	
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.	

## Data

KDa	М	R
135 100 75 65 45	101	
35		
25		
15	-	ī

> 95 % as determined by reducing SDS-PAGE.

## Background

Angiopoietin-1 (Ang-1) is a secreted glycoprotein that plays a critical role in the development and maintenance of the vascular system. The angiopoietin (ANGPT)-TIE2/TEK signaling pathway is essential for blood and lymphatic vascular homeostasis. ANGPT1 is a potent TIE2 activator, whereas ANGPT2 functions as a context-dependent agonist/ antagonist. Angiopoietin-1 can protect against pulmonary arterial hypertension, reduce the extent of fibrosis and remodeling in infarcted diabetic myocardium, and enhance tumor progression and metastasis.