

## Recombinant Mouse MEK4/MAP2K4 Protein (His & GST Tag)

**Catalog Number:** PKSM040567

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	Baculovirus-Insect Cells-derived Mouse MEK4/MAP2K4 protein Ala 2-Asp 397, with an N-terminal His & GST
<b>Mol_Mass</b>	72.0 kDa
<b>Accession</b>	P47809
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Dual specificity mitogen-activated protein kinase kinase 4, also known as MAP kinase kinase 4, MAPKK4, JNK-activating kinase 1, MAPK/ERK kinase 4, SAPK/ERK kinase 1, c-Jun N-terminal kinase kinase 1, JNKK and MAP2K4, is a protein which belongs to the protein kinase superfamily, STE Ser/Thr protein kinase family and MAP kinase kinase subfamily. MAP2K4 / JNKK1 is a protein kinase which is a direct activator of MAP kinases in response to various environmental stresses or mitogenic stimuli. MAP2K4 / JNKK1 has been shown to activate MAPK8 / JNK1, MAPK9 / JNK2, and MAPK14 / p38, but not MAPK1 / ERK2 or MAPK3 / ERK1. MAP2K4 / JNKK1 is phosphorylated, and thus activated by MAP3K1 / MEKK. The stress-activated protein kinase (SAPK) pathways represent phosphorylation cascades that convey pro-apoptotic signals. The mitogen-activated protein kinase kinase (MAPKK) homolog MAP2K4 (MKK4, SEK, JNKK1) is a centrally-placed mediator of the SAPK pathways.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V3.5