

Recombinant Mouse PLA2G2E Protein (His Tag)

Catalog Number: PKSM040548

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse PLA2G2E protein Met 1-Cys 142, with an C-terminal His
Calculated MW	15.5 kDa
Observed MW	18 kDa
Accession	Q9QUL3
Bio-activity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, 10% glycerol, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Group IIE secretory phospholipase A2, also known as GIIE sPLA2, sPLA2-IIE, Phosphatidylcholine 2-acylhydrolase 2E and PLA2G2E is a secreted protein which belongs to the phospholipase A2 family. Mammalian secretory phospholipase A2s (sPLA2s) form a family of structurally related enzymes that are involved in a variety of physiological and pathological processes via the release of arachidonic acid from membrane phospholipids or the binding to specific membrane receptors. Phospholipases A2 / PLA2 are enzymes that release fatty acids from the second carbon group of glycerol. This particular phospholipase specifically recognizes the sn-2 acyl bond of phospholipids and catalytically hydrolyzes the bond releasing arachidonic acid and lysophospholipids. Phospholipases A2 / PLA2 are commonly found in mammalian tissues as well as insect and snake venom. Venom from both snakes and insects is largely composed of melittin, which is a stimulant of Phospholipases A2 / PLA2. Due to the increased presence and activity of Phospholipases A2 / PLA2 resulting from a snake or insect bite, arachidonic acid is released from the phospholipid membrane disproportionately. As a result, inflammation and pain occur at the site. PLA2G2E catalyzes the calcium-dependent hydrolysis of the 2-acyl groups in 3-sn-phosphoglycerides. Has a preference for arachidonic-containing phospholipids.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017