

## Recombinant Human COL2A1 Protein (aa 1242-1487, His Tag)

**Catalog Number:** PKSH031001

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human COL2A1 protein Asp 1242-Leu 1487, with an N-terminal His
<b>Calculated MW</b>	29.8 kDa
<b>Observed MW</b>	37 kDa
<b>Accession</b>	P02458-2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 92 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 92 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

COL2A1 is the alpha-1 chain of type II collagen which is a fibrillar collagen found in cartilage and the vitreous humor of the eye. Mutations in this protein are associated with achondrogenesis, chondrodysplasia, early onset familial osteoarthritis, SED congenita, Langer-Saldino achondrogenesis, Kniest dysplasia, Stickler syndrome type I, and spondyloepimetaphyseal dysplasia Strudwick type. In addition, defects in processing chondrocalcin, a calcium binding protein that is the C-propeptide of this collagen molecule, are also associated with chondrodysplasia. There are two transcripts identified for this gene. Type II collagen is specific for cartilaginous tissues. Thus COL2A1 is essential for the normal embryonic development of the skeleton, for linear growth and for the ability of cartilage to resist compressive forces. The regulation of COL2A1, likely results from a balance of both positive and negative proteins. The inhibition of COL2A1 transcription following treatment of chick sternal chondrocytes with growth factors was accompanied by increased EF1 expression. Overexpression of EF1 in differentiated chondrocytes resulted in decreased expression of a reporter construct containing a collagen II promoter/enhancer insert.