A Reliable Research Partner in Life Science and Medicine

(KO Validated) FGF2 Polyclonal Antibody

catalog number: E-AB-93292

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide of human FGF2

Host Rabbit Isotype IgG


Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Recommended Dilution Applications

1:500-1:2000 WB

Data





Western blot analysis of Mouse ovary using FGF2 Polyclonal Western blot analysis of extracts from wild type(WT) and Antibody at 1:2500 dilution.

Observed-MW:18 kDa/22 kDa/24 kDa Calculated-MW:17 kDa/21 kDa/22 kDa/30 kDa FGF2 knockout (KO) HeLa cells using FGF2 Polyclonal

Antibody at 1:2000 dilution.

Observed-MW:18 kDa/22 kDa/24 kDa Calculated-MW:17 kDa/21 kDa/22 kDa/30 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.

For Research Use Only

Fax: 1-832-243-6017