

Recombinant Mouse CD27/TNFRSF7 Protein(Fc Tag)

Catalog Number: PDMM100150

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	Mammalian-derived Mouse CD27/TNFRSF7 proteins Pro24-Arg182, with an C-terminal Fc
Calculated MW	42.4 kDa
Observed MW	60 kDa
Accession	P41272
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse CD27/TNFRSF7 proteins ,
2µg/lane of Recombinant Mouse CD27/TNFRSF7 proteins
was resolved with SDS-PAGE under reducing conditions ,
showing bands at 60 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7

CD27, also known as TNFRSF7, is a member of the TNF-receptor superfamily limited to cells of the lymphoid lineage, and exists as both a dimeric glycoprotein on the cell surface and as a soluble protein in serum. As a type I transmembrane glycoprotein of about 55 kDa existing as disulfide-linked homodimer, CD27 has been shown to play roles in lymphoid proliferation, differentiation, and apoptosis. It has an important role in the generation of T cell immunity and is an robust marker for normal memory B cells. It is a T and B cell co-stimulatory molecule, the activity of CD27 is governed by its TNF-like ligand CD70 on lymphocytes and dendritic cells. The CD27-CD70 interaction is required for Th1 generation responses to differentiation signals and long-term maintenance of T cell immunity, and meanwhile, plays a key role in regulating B-cell differentiation, activation and immunoglobulin synthesis.