# **Elabscience**®


## Recombinant Human Pro-ADM protein (GST Tag)

### Catalog Number: PDEH100786

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description    |  |
|----------------|--|
| Species        | Human  |
| Source         | E.coli-derived Human Pro-ADM protein Arg22-Tyr146, with an N-terminal GST  |
| Calculated MW  | 38.6 kDa   |
| Observed MW    | 30 kDa   |
| Accession      | P35318   |
| Bio-activity   | Not validated for activity   |
| Properties     |  |
| Purity         | >95% as determined by reducing SDS-PAGE.   |
| Endotoxin      | < 10 EU/mg of the protein as determined by the LAL method  |
| Storage        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80   |
|                | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20$ °C for 3 months. |
| Shipping       | This product is provided as lyophilized powder which is shipped with ice packs.  |
| Formulation    | Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%   |
|                | Mannitol.  |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of   |
|                | 0.5 mg/mL. Concentration is measured by UV-Vis.  |

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

AM and PAMP are potent hypotensive and vasodilatator agents. Numerous actions have been reported most related to the physiologic control of fluid and electrolyte homeostasis. In the kidney, am is diuretic and natriuretic, and both am and pamp inhibit aldosterone secretion by direct adrenal actions. In pituitary gland, both peptides at physiologically relevant doses inhibit basal ACTH secretion. Both peptides appear to act in brain and pituitary gland to facilitate the loss of plasma volume, actions which complement their hypotensive effects in blood vessels.