



A Reliable Research Partner in Life Science and Medicine

Human MMP-1 Antibody Pair Set

Catalog No. E-KAB-0053 Applications ELISA

Synonyms MMP1, CLG, CLGN, Interstitial Collagenase, Fibroblast Collagenase

Kit components & Storage

Title	Specifications	Storage
Human MMP-1 Capture Antibody	1 vial, 100 μ g	Store at -20℃. Avoid freeze/thaw
		cycles.
Human MMP-1 Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze/thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0053)		
		Human MMP-1 Capture Antibody	Human MMP-1 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human MMP-1 protein	Recombinant Human MMP-1 protein	
Information	Swissprot	P03956		
Product details	Reactivity	Human	Human	
	Host	Goat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol, pH 7.4	protective protein; 50% glycerol; pH	
			7.4	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Human MMP-1 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human MMP-1 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human MMP-1 Capture Antibody	
Capture			Onlical Density
ELISA	1:1000-1:10000	Human MMP-1 Detection Antibody	Optical Christian
Detection		(Biotin)	,
			100 100 15

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes a secreted enzyme which breaks down the interstitial collagens, types I, II, and III. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. Alternative splicing results in multiple transcript variants.

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u> Fax: 1-832-243-6017