

## Human MMP-1 Antibody Pair SetSet

**Catalog No.** E-KAB-0053 **Applications** ELISA  
**Synonyms** MMP1, CLG, CLGN, Interstitial Collagenase, Fibroblast Collagenase

### Kit components & Storage

Title	Specifications	Storage
Human MMP-1 Capture Antibody	1 vial, 100 µg	Store at -20℃. Avoid freeze/thaw cycles.
Human MMP-1 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20℃. Avoid freeze/thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Product Information

Items		Characteristic (E-KAB-0053)	
		Human MMP-1 Capture Antibody	Human MMP-1 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human MMP-1 protein	Recombinant Human MMP-1 protein
	Swissprot	P03956	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human MMP-1 in ELISAs.	

### For Research Use Only

## Applications

### Human MMP-1 Sandwich ELISA Assay

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human MMP-1 Capture Antibody	
ELISA Detection	1:1000-1:10000	Human MMP-1 Detection Antibody (Biotin)	

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes a secreted enzyme which breaks down the interstitial collagens, types I, II, and III. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. Alternative splicing results in multiple transcript variants.

## For Research Use Only