

P-Selectin/CD62P Polyclonal Antibody(Capture/Detector)

catalog number: AN003860P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Mouse |
| Immunogen | Recombinant Mouse P-Selectin/CD62P Protein expressed by Mammalian |
| Host | Rabbit |
| Isotype | Rabbit IgG |
| Purification | Antigen Affinity Purification |
| Conjugation | Unconjugated |
| Buffer | Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300. |

Applications

Recommended Dilution

| | |
|-----------------------|---------------|
| ELISA Capture | 2-8 µg/mL |
| ELISA Detector | 0.1-0.4 µg/mL |

Data



Sandwich ELISA-Recombinant Mouse P-Selectin/CD62P Protein standard curve. Background subtracted standard curve using Anti-P-Selectin/CD62P antibody(AN003860P) (Capture), Anti-P-Selectin/CD62P antibody(AN003860P) (Detector). The reference range value is 0.16-10 ng/mL for mouse.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

For Research Use Only

Mouse P-Selectin (GMP-140, LECAM-3, PADGEM, CD62P), a member of the Selectin family, is a cell surface glycoprotein expressed by activated platelets and endothelial cells. P-Selectin is translocated to the cell surface within minutes, from alpha granules of platelets or Weibel-Palade bodies of endothelial cells, following stimulation with thrombin, histamine, PMA or peroxides. P-Selectin binds to a 106 kDa protein present on myeloid cells, neutrophils, monocytes and lymphocytes, termed PSGL-1 (P-Selectin glycoprotein ligand-1). P-Selectin plays a role in the adhesion of leukocytes and neutrophils to the endothelium. Acting in cooperation with L-Selectin, P-Selectin mediates the initial interaction of circulating leukocytes with endothelial cells that produces a characteristic 'rolling' of the leukocytes on the endothelium. This initial interaction is followed by a stronger interaction involving E-Selectin, and later ICAM-1 and VCAM-1, that leads eventually to extravasation of the white blood cell through the blood vessel wall into the extracellular matrix tissue. Mouse P-Selectin cDNA encodes a 768 amino acid (aa) residue type I transmembrane protein with a 41 aa signal peptide, a 668 aa extracellular domain, a transmembrane domain and a short (35 aa) cytoplasmic domain. The extracellular domain has an NH2-terminal C-type lectin domain and an EGF-like domain followed by a series of complement factor A repeat homology domains. The extracellular domains of human and mouse P-Selectin share approximately 73% sequence homology.