

## CD74 Monoclonal Antibody

catalog number: **AN200025P**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CD74 protein
<b>Host</b>	Mouse
<b>Isotype</b>	IgG1
<b>Clone</b>	A869
<b>Purification</b>	Protein A
<b>Buffer</b>	0.2 µm filtered solution in PBS


### Applications Recommended Dilution

<b>IHC-P</b>	1:500-1:2000
<b>FCM</b>	1:25-1:100


### Data




Flow cytometric analysis of Human CD74 expression on human whole blood lymphocytes. Cells were stained with purified anti-Human CD74, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of viable lymphocytes.



Immunohistochemistry of paraffin-embedded human tonsil using CD74 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded human colon carcinoma using CD74 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded human appendix using CD74 Monoclonal Antibody at dilution of 1:1000.

### Preparation & Storage

<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Shipping</b>	Ice bag

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.2

CD74 is a type II transmembrane glycoprotein also known as MHC class II associated invariant chain, invariant chain, Ii, MHC class II chaperone, and MIF receptor. CD74 exists in four isoforms with molecular masses of 33, 35, 41, and 43 kD, depending on genetic splicing. CD74 is primarily expressed on antigen presenting cells, including B cells, monocytes/macrophages, dendritic cells, and Langerhans cells. It is also expressed by activated T cells and activated endothelial and epithelial cells as well as carcinomas of lung, renal, gastric and thymic origin. The primary function of CD74 is intracellular sorting of MHC class II molecules and regulation of exogenous peptide loading onto MHC class II. It is also involved in the modulation of B cell differentiation and positive selection of CD4+ T cells. It has been reported that CD74 binds MIF (macrophage migration inhibitory factor) and signals through CD44 to regulate innate and adaptive immunity. It is also reported that H. pylori infection occurs through urease B binding of CD74 on gastric epithelial cells, inducing gastric epithelial cell apoptosis, NF-κB activation, and IL-8 production.