

## Recombinant Mouse IL18BP Protein (Fc Tag)

**Catalog Number:** PKSM041077

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse IL18BP protein Thr29-Ala193, with an C-terminal Fc
<b>Calculated MW</b>	45.1 kDa
<b>Observed MW</b>	60-90 kDa
<b>Accession</b>	Q9Z0M9
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Interleukin 18 binding protein (IL-18BP) is a physiological inhibitor that acts through binding to the receptor-binding site of IL-18. IL-18 stimulates INF- $\gamma$ , which then stimulates 18BP production via NF- $\kappa$ B. The interaction between IL-18 and IL-18BP has a significant role in the inflammation process. IL-18 BPs have four isoforms, a, b, c and d, which are spliced by different ways. The IL-18 BP isoforms a and c each contain one immunoglobulin (Ig)-like C2-type domain which is essential to the binding and neutralizing properties of the binding proteins. The IL-18 BP isoforms b and d lack a complete Ig domain. The expression of IL-18 and the IL-18BP are identified in immune tissues such as the spleen, but also in nonimmune cells including keratinocytes.

### For Research Use Only