

## Purified Anti-Mouse CD205 Antibody[NLDC-145], Functional Grade

catalog number: AN008440

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |                                                                                       |
|---------------------|---------------------------------------------------------------------------------------|
| <b>Reactivity</b>   | Mouse                                                                                 |
| <b>Immunogen</b>    | Recombinant Mouse CD205 protein                                                       |
| <b>Host</b>         | Rat                                                                                   |
| <b>Isotype</b>      | Rat Wl, also known as Wistar (outbred) IgG2a, κ                                       |
| <b>Clone</b>        | NLDC-145                                                                              |
| <b>Purification</b> | >98%, Protein A/G purified                                                            |
| <b>Buffer</b>       | Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method. |

### Applications


### Recommended Dilution

**FCM** ≤ 0.2 µg per million cells in 100 µL volume

### Data



A20 were stained with 0.2 µg Purified Anti-Mouse CD205 Antibody[NLDC-145], Functional Grade (Right) and 0.2 µg Rat IgG2a, κ Isotype Control(Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.



A20 were stained with 0.2 µg Purified Anti-Mouse CD205 Antibody[NLDC-145], Functional Grade (Right) and 0.2 µg Rat IgG2a, κ Isotype Control(Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.

### Preparation & Storage

|                 |                                                                                                                                                                                                    |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions. |
| <b>Shipping</b> | Ice bag                                                                                                                                                                                            |

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.4

CD205, also known as DEC-205, is a 205 kD integral membrane protein homologous to the macrophage mannose receptor. It is a type I cell surface protein that belong to the C-type lectin family. CD205 is expressed at high levels by dendritic cells and thymic epithelial cells. It is also expressed by a number of other cell types, such as B lymphocytes, macrophages, Langerhans cells, bone marrow stromal cells, granulocytes, epithelial cells of pulmonary airways, and the capillaries of the brain. CD205 is a novel endocytic receptor used by dendritic cells and thymic epithelial cells to direct captured antigens from the extracellular space to specialized antigen processing. It mediates antigen uptake and presentation and cross-presentation to T cells. It has been reported that CD205 acts as a recognition receptor for dying cells, potentially provides an important pathway for the uptake of self-antigen in the intrathymic environment, and is involved in peripheral tolerance. Antibody-mediated antigen-targeting via the DEC-205 receptor increases the efficiency of vaccination for T cell immunity.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.