ATXN3 Polyclonal Antibody

catalog number: E-AB-18525



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--|--|
| Reactivity | Human |
| Immunogen | Fusion protein of human ATXN3 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:50-1:300 |
| Data | |
| Immunohistochemistry of paraffin-embedded Human liver cancer tissue using ATXN3 Polyclonal Antibody at dilution of 1:100(×200)Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using ATXN3 Polyclonal Antibody at dilution of 1:100(×200) | |
| Preparation & Storage | |
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |
| Background | |
| Maahada Jaganh digagaa alaa kugum aganingaamhallanatayig 2 igan aytagamal daminant nayralagia digardar. Tha | |

Machado-Joseph disease, also known as spinocerebellar ataxia-3, is an autosomal dominant neurologic disorder. The protein encoded by this gene contains (CAG)n repeats in the coding region, and the expansion of these repeats from the normal 12-44 to 52-86 is one cause of Machado-Joseph disease. There is a negative correlation between the age of onset and CAG repeat numbers. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.

For Research Use Only