

Recombinant Human R-Cadherin/CDH4 Protein (His Tag)

Catalog Number: PKSH031741

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human R-Cadherin/CDH4 protein Met 1-Ala 734, with an C-terminal His
Mol_Mass	80.0 kDa
Accession	NP_001785.2
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat brain glial cells. Immobilized CAD4 (0.8 µg/ml, 100 µl/well) will mediate > 20% C6 cell adhesion.

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

The cadherin superfamily is a large family that engage in both homo- and heterotypic, calcium-dependent, cell-cell adhesion events, and can be divided into at least four subfamilies based on the extracellular (EC) regions and cytoplasmic domains, that is: classical cadherins, desmosomal cadherins, protocadherins, and cadherin-like molecules. Human cadherin 4, type 1, R-cadherin (retinal), also known as CDH4, CAD4 and RCAD, is a classical cadherin from the cadherin superfamily. It is a calcium-dependent adhesion molecule and a type I transmembrane glycoprotein composed of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. CDH4 is thought to play an important role during brain segmentation and neuronal outgrowth, and also exerts critical actions in kidney and muscle development. CDH4 is expressed in vascular smooth muscle, pancreatic β -cells, thyroid follicular cells, sensory neurons of the dorsal root ganglia, and, possibly, astrocytes and endothelium of the retina. As a classic cadherin, CDH4 forms both homodimers and heterodimers with N-cadherin. The extracellular region of human CDH4 is 96% aa identical to that of mouse CDH4.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017