

A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410]

catalog number: E-AB-F1016A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Host Mouse

 $\begin{tabular}{lll} \textbf{Isotype} & Mouse IgGl, \kappa \\ \textbf{Clone} & MIAP410 \\ \textbf{Conjugation} & Unconjugated \\ \end{tabular}$


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze

to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



Rat splenocytes were stained with 0.2 μg Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracelluluar domain, a five membrane-spanning segment and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRPα, CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.

For Research Use Only