RPL13A Polyclonal Antibody

catalog number: E-AB-18938



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Fusion protein of human RPL13A

Host Rabbit Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilut	ion

WB 1:1000-1:5000 **IHC** 1:50-1:300


Data



Western blot analysis of LOVO Hela A549 and HT29 cell lysates using RPL13A Polyclonal Antibody at dilution of 1:1600

Immunohistochemistry of paraffin-embedded Human tonsil tissue using RPL13A Polyclonal Antibody at dilution of 1:95(×200)

Observed-MV:Refer to figures



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using RPL13A Polyclonal Antibody at dilution of $1.95(\times 200)$

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

RPL13A Polyclonal Antibody

catalog number: E-AB-18938



Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a member of the L13P family of ribosomal proteins that is a component of the 60S subunit. The encoded protein also plays a role in the repression of inflammatory genes as a component of the IFN-gamma-activated inhibitor of translation (GAIT) complex. This gene is co-transcribed with the small nucleolar RNA genes U32, U33, U34, and U35, which are located in the second, fourth, fifth, and sixth introns, respectively. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.