

## Recombinant Rat IL-22R/IL22RA1 Protein (ECD, Fc Tag)

**Catalog Number:** PKSR030161

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat IL-22R/IL22RA1 protein Met1-Ala228, with an C-terminal hFc
<b>Calculated MW</b>	50.8 kDa
<b>Accession</b>	NP_001178798.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

IL-22R belongs to the type II cytokine receptor family. It contains 2 fibronectin type-III domains and is expressed in colon, liver, lung, pancreas and kidney. IL-22R also can be expressed in keratinocytes of normal skin as well as in psoriatic skin lesion. Overexpression of IL-22R can be detected in synovial fluid cells from rheumatoid arthritis and spondyloarthropathy patients. IL-22R is a component of the receptor for IL20, IL22 and IL24. The component of IL-22R formed by IL22RA1 and IL10RB enables IL22 signaling via JAK/STAT pathways. IL22 also induces activation of MAPK1/MAPK3 and Akt kinases pathways. Component of one of the receptor for IL20 and IL24 formed by IL22RA1 and IL20RB also signaling through STATs activation. IL-22R mediates IL24 antiangiogenic activity as well as IL24 inhibitory effect on endothelial cell tube formation and differentiation.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017