



A Reliable Research Partner in Life Science and Medicine

# **Human FETUA Antibody Pair Set**

Catalog No. E-KAB-0178 Applications ELISA

Synonyms AHSG, A2HS, AHS, HSGA, alpha-2-HS-glycoprotein

## **Kit components & Storage**

Title	Specifications	Storage
Human FETUA Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human FETUA Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0178)		
		Human FETUA Capture Antibody	Human FETUA Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human FETUA protein	Recombinant Human FETUA protein	
Information	Swissprot	P02765		
Product details	Reactivity	Human	Human	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Human FETUA in ELISAs.	·	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human FETUA Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human FETUA Capture Antibody	
Capture			Aig
ELISA Detection	1:1000-1:10000	Human FETUA Detection Antibody (Biotin)	O 0.01 100 1000 10000 Human FETUA concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

Alpha2-HS glycoprotein (AHSG), a glycoprotein present in the serum, is synthesized by hepatocytes. The AHSG molecule consists of two polypeptide chains, which are both cleaved from a proprotein encoded from a single mRNA. It is involved in several functions, such as endocytosis, brain development and the formation of bone tissue. The protein is commonly present in the cortical plate of the immature cerebral cortex and bone marrow hemopoietic matrix, and it has therefore been postulated that it participates in the development of the tissues. However, its exact significance is still obscure.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>