

A Reliable Research Partner in Life Science and Medicine

PE Anti-Rat CD45 Antibody[OX-1]

Catalog Number: E-AB-F1227UD

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. OX-1

PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793D] Isotype Control

Conjugation

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

> check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

> reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10⁶ cells

in 100 µL volume].

Data



Rat splenocytes are stained with PE Anti-Rat CD45 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. **Storage**

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names LCALy-5T200;Leukocyte common antigen;Ptprc;Receptor-type tyrosine-protein

phosphatase C

Uniprot ID P04157 Gene ID 19265

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD45 is a 180-220 kD protein also known as leukocyte common antigen (LCA). It is a protein tyrosine phosphatase with multiple isoforms differing as a result of alternative splicing of the extracellular domain and glycosylation. CD45 is expressed on all hematopoietic cells except erythrocytes and platelets; isoform expression depends on cell type, activation state, and cell maturation. CD45 functions in signal transduction through T and B cell antigen receptors. CD45 has been shown to interact with various proteins including galectin-1, CD2, CD3, and CD4. The OX-1 antibody has been shown to partially inhibit NK cell-mediated lysis of syngeneic tumor cells in vitro.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com Email:techsupport@elabscience.com