



A Reliable Research Partner in Life Science and Medicine

Recombinant Histone H4 (Acetyl Lys8) Monoclonal Antibody

catalog number: AN301277L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide corresponding to residues around (Lys8) of Human Acetyl-

Histone H4

 Host
 Rabbit

 Isotype
 IgG,κ

 Clone
 B1040


 Purification
 Protein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

IHC 1:500-1:2000

Data



Immunohistochemistry of paraffin-embedded human squamous cell lung carcinoma using Recombinant Histone H4 (Acetyl Lys8) Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1