TP63 Polyclonal Antibody

catalog number: E-AB-18722



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Reactivity Immunogen | Human;Mouse;Rat |
|--|---|
| Immunogen | |
| - | Fusion protein of human TP63 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:50-1:200 |
| Data | |
| De se | |
| • • | paraffin-embedded Human lung blyclonal Antibody at dilution of |
| cancer tissue using TP63 Pc | paraffin-embedded Human lung olyclonal Antibody at dilution of 55(×200) |
| cancer tissue using TP63 Pc 1:6 | blyclonal Antibody at dilution of |
| cancer tissue using TP63 Pc | blyclonal Antibody at dilution of |
| cancer tissue using TP63 Pc 1:6 Preparation & Storage | olyclonal Antibody at dilution of 55(×200) |
| cancer tissue using TP63 Pc 1:6 Preparation & Storage Storage | blyclonal Antibody at dilution of 55(×200) Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| cancer tissue using TP63 Pc 1:6 Preparation & Storage Storage | blyclonal Antibody at dilution of (×200) Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the |

p53 and p73 genes. It is involved in malignancy acquisition and maintenance of cells. Unlike p53,the p63 gene encodes multiple isotypes with remarkably divergent abilities to transactivate p53 reporter genes and induce apoptosis. TP63 acts as a sequence specific DNA binding transcriptional activator or repressor. TP63 has 12 isoforms with MW 40kd(P40), 50kd(P60),63kd(P63) and 73kd(P73L). It is Nuclear stain.

For Research Use Only